

High Prevalence of GB Virus C Strains Genetically Related to Strains With Asian Origin in Nicaraguan Hemophiliacs

Mario A. González-Pérez,^{1,3} Heléne Norder,^{1*} Åsa Bergström,¹ Ernesto López,² Kirsten A. Visoná,³ and Lars O. Magnius¹

¹Department of Virology, Swedish Institute for Infectious Disease Control, Stockholm, Sweden

²Red Cross Blood Program, Red Cross of Nicaragua, Managua, Nicaragua

³Louisiana State University–International Center for Medical Research and Training, San José, Costa Rica

The presence of hepatitis GB virus C (GBV-C), also known as hepatitis G virus (HGV), and hepatitis C virus (HCV) were investigated in sera from 45 hemophiliacs from nine locations in Nicaragua using a nested polymerase chain reaction (PCR). Primers used to detect GBV-C and HCV derived from the helicase region and 5'UTR, respectively. Seventeen (38%) patients were positive for GBV-C RNA in serum by PCR. Twelve (27%) patients were positive for HCV RNA by PCR. Six (13%) of these were coinfecting with GBV-C. Anti-HCV was detected in all the 12 HCV RNA positive hemophiliacs and in another 14 (31%) individuals, in whom GBV-C RNA was found in 2. Ten patients (22%) lacked markers for both GBV-C and HCV. The mean age of the patients positive for GBV-C but negative for HCV by PCR was significantly lower than for those negative for GBV-C but positive for HCV by PCR ($P < 0.05$; Student's *t*-test), indicating that the risk for this group of hemophiliacs to acquire GBV-C infection is higher as compared to the risk of acquiring HCV infection. Eleven GBV-C strains were sequenced in the 5'UTR. Sequence comparison to previously published GBV-C strains revealed that all 11 strains were more similar to Asian strains than to strains of European and African origin. Sequences in the NS5-B region were available for 8 HCV strains, all of which were found to belong to genotype 1a. The similarity of the Nicaraguan GBV-C strains to strains from Asia indicates that the GBV-C strains in the region presumably have an Amerindian origin. It is also considered that the HTLV II strains in the New World aboriginal populations are ancient and brought there by the ancestral Amerindian populations from Asia. Further, the genotype F of hepatitis B virus, known to represent the strains in populations with Amerindian background, predominates in Central American populations with Hispanic background. It re-

mains to be clarified why Amerindian strains of GBV-C as well as of HBV predominate also in populations with mixed ethnic background in Central America. *J. Med. Virol.* 52:149–155, 1997. © Wiley-Liss, Inc.

KEY WORDS: GBV-C; sequences; hemophiliacs; Amerindians; Central America

INTRODUCTION

A new virus designated GBV-C implicated in the etiology of viral hepatitis was described recently [Simons et al., 1995]. This virus is a positive single-stranded RNA virus of the Flaviviridae family more related to the previously discovered GBV-A and GBV-B and hepatitis C virus (HCV) than to pestiviruses and true flaviviruses, e.g., yellow fever virus, dengue viruses, and others [Simons et al., 1995]. Another recently described virus designated as hepatitis G virus (HGV) [Linnen et al., 1996] shares 95% homology at the amino acid level with GBV-C and therefore just represents another isolate of this virus [Zuckerman, 1996]. In contrast to HCV, for which 11 different genotypes are described with up to 30% variation based on the complete open reading frame [Choo et al., 1991; Okamoto et al., 1992; Tokita et al., 1994a; 1994b; 1995], GBV-C is a highly conserved virus with a divergency at the amino acid level of the polyprotein of only 4% between strains from the United States and Europe represented by HGV and African strains represented by GBV-C. However, based on phylogenetic analysis of the 5'-UTR,

Contract grant sponsor: Swedish Medical Research Council; contract grant number K96-16P-11019-03 A.

*Correspondence to: Heléne Norder, Virological Department, Swedish Institute for Infectious Disease Control, S105 21 Stockholm, Sweden.

Accepted 13 January 1997

there are 3 different GBV-C clusters representing strains of varying geographical origin, tentatively designated as genotypes 1, 2, and 3 [Muerhoff et al., 1996].

The presence of divergent genomes of several different hepatitis viruses, e.g., hepatitis B virus [Naumann et al., 1993], hepatitis E virus [Huang et al., 1992], and hepatitis D virus [Casey et al., 1993], confined to certain ethnic groups in America, directs the attention of investigators to search for divergent viral genotypes in the area. So far, only scant information is available regarding HCV genotypes in Central America [Yun et al., 1996] and no information is available regarding the genetic diversity of GBV-C. This study was conducted in order to determine the presence of GBV-C and the genotypes of HCV in a group of hemophiliac patients at high risk of contracting viral hepatitis in Nicaragua.

MATERIALS AND METHODS

Study group

Serum samples from 45 hemophiliac patients (age range: 1–31 years, mean \pm SD: 12.5 \pm 7.1) from nine different locations in Nicaragua, were obtained, aliquotted, and shipped on dry ice to the Department of Virology at the Swedish Institute for Infectious Disease Control, Stockholm, Sweden.

Serological Methods

Sera were tested for the presence of anti-HCV using a second generation anti-HCV test (Abbott Laboratories, Chicago, IL) at Red Cross Blood Bank, Managua, Nicaragua. Positive samples were analysed with the recombinant immunoblot assay, RIBA HCV 3.0 (Chiron, Emeryville, CA) at the Swedish Institute for Infectious Disease Control.

RNA Extraction and cDNA Synthesis

RNA was extracted from serum according to Garson et al. [1990] with minor modifications. Briefly, 100 μ l serum was precipitated with 10% polyethylene glycol in 0.03M NaCl for 40 minutes on ice, the tubes were centrifuged, and the precipitate was dissolved in 125 μ l of a buffer containing 5mM NaCl, 10mM Tris-HCl pH 8.3, 1mM EDTA, pH 7.0, and 25ng tRNA; 40 μ l of a second buffer consisting of 5% sodium dodecyl sulphate, 25mM EDTA pH 7.0, and 250mM potassium acetate was added and the tubes were shaken vigorously at room temperature for 5 minutes. After phenol and phenol/chloroform extraction, the nucleic acids were ethanol precipitated. The RNA was dissolved in 20 μ l diethyl-pyrocabonate treated water and 5 μ l dissolved RNA was used for cDNA synthesis using 1 U Superscript II reverse transcriptase (Gibco, BRL) and 1 nmole of random hexamer primers (Boehringer Mannheim).

GBV-C PCR and Sequencing

The NS3 region of the cDNA was amplified by PCR using primers GB1 (5'-GCCAACCCKAGGCAGAT-GCT-3') at position 4035–4054 of the GBV-C genome according to Simons et al. [1995] and GB2 (5'-GAYAG-

CGCGTCTGTGMCACA-3') at position 4433–4452. The product obtained was nested with primers GBVC-s1/GBVC-a1 according to Leary et al. [1996]. The 5'-UTR was amplified by PCR using primers S1/GBVCE1wb2 according to Muerhoff et al. [1996]. Five μ l of the product was further amplified with the primers S1/4R [Muerhoff et al., 1996]. The PCR product was purified with Wizard PCR preps DNA purification system (Promega) and 0.5 pmoles were sequenced according to Casanova et al. [1990] in both directions with 5 pmoles of the primers S1 and 4R.

HCV PCR and Sequencing

HCV RNA was detected in the samples by amplification of the cDNA in the 5'-UTR with the primer pair NCR1/NCR2 [Garson et al., 1990]. Five μ l of the obtained product was further amplified with primers NCR3/NCR4 [Garson et al., 1990].

The NS5-B region of the cDNA was amplified by PCR using primers hep120 (5'-TGCGCGACBGABACRTT-KGAGGA-3') at position 8702–8724 of the HCV genome according to Choo et al. [1991] and primer hep101 (5'-ATACCCGCTGCTTTGACTC-3') at position 8258–8276. The product obtained was amplified further with hep101b as hep 101 with 5'-biotinylation and hep 105 (5'-ATACCTAGTCATAGCCTCCGTGA-3') at position 8616–8638. The PCR product was purified with Wizard PCR preps DNA purification system (Promega) and used for solid-phase DNA sequencing. One pmol of the purified product was immobilized on magnetic beads (Dynabeads M280-streptavidin; Dynal AS) and subsequently denatured to obtain single stranded DNA [Leitner et al., 1995]. The sequencing reaction was carried out for both strands using the reagents provided in the Sequenase Version 2.0 DNA Sequencing Kit (USB Cleveland, OH), [³⁵S]dATP, 2 units of Sequenase (USB), and 5 pmoles of the primers used in the PCR as sequencing primers. The sequenced fragments were electrophoresed on 6% polyacrylamide gels and autoradiographed.

Tree Constructions

Of the sequences obtained, 375 bp were aligned with the corresponding part of the genome of 49 GBV-C sequences available in EMBL GeneBank (Table I) and with GBV-C sequences in the sera from three Swedish drug addicts. Dendrograms were created with the programs DNADIST together with NEIGHBOUR (UPGMA) in the PHYLIP package, version 3.53 [Felsenstein, 1993]. SEQBOOT was used to bootstrap data, in which 100 data sets were analysed, and CONSENSE was used to compute consensus trees. For visualization and graphical editing of the trees, the program TREETOOL, version 1.0, in GDE was used.

RESULTS

Results are summarized in Tables II and III. GBV-C RNA was detected by nested-PCR using primers in the NS3 region in seventeen (38%) of the hemophiliacs (age range: 2–21; mean \pm SD: 12.3 \pm 6.4). Twenty-eight

TABLE I. Designation and GenBank Accession Numbers of the GBV-C Sequences Used in the Comparison

Designation	Genotype	Origin	Accession no.	Reference
Isolate 23	1a	West Africa	U59540	Muerhoff et al., 1996
Isolate 24	1a	West Africa	U59541	Muerhoff et al., 1996
Isolate 25	1a	West Africa	U59542	Muerhoff et al., 1996
Isolate 26	1a	West Africa	U59543	Muerhoff et al., 1996
Isolate 27	1a	West Africa	U59544	Muerhoff et al., 1996
Isolate 28	1a	West Africa	U59545	Muerhoff et al., 1996
Isolate 29	1a	West Africa	U59546	Muerhoff et al., 1996
GBV-C	1b	West Africa	U36330	Leary et al., 1996
Isolate 31	1b	West Africa	U59547	Muerhoff et al., 1996
Isolate 32	1b	West Africa	U59548	Muerhoff et al., 1996
Isolate 33	1b	West Africa	U59549	Muerhoff et al., 1996
Isolate 34	1b	West Africa	U59550	Muerhoff et al., 1996
Isolate 35	1b	West Africa	U59551	Muerhoff et al., 1996
Isolate 36	1b	West Africa	U59552	Muerhoff et al., 1996
Isolate 37	1b	West Africa	U59553	Muerhoff et al., 1996
Isolate 38	1b	West Africa	U59554	Muerhoff et al., 1996
Isolate 39	1b	West Africa	U59555	Muerhoff et al., 1996
Isolate 40	1b	West Africa	U59556	Muerhoff et al., 1996
Isolate 41	1b	West Africa	U59557	Muerhoff et al., 1996
Isolate 42	1b	West Africa	U59558	Muerhoff et al., 1996
PNF2161	2a	USA	U44402	Linnen et al., 1996
Isolate 1	2a	USA	U59518	Muerhoff et al., 1996
Isolate 2	2a	USA	U59519	Muerhoff et al., 1996
Isolate 3	2a	Europe	U59520	Muerhoff et al., 1996
Isolate 4	2a	Europe	U59521	Muerhoff et al., 1996
Isolate 5	2a	Europe	U59522	Muerhoff et al., 1996
Isolate 6	2a	USA	U59523	Muerhoff et al., 1996
Isolate 7	2a	USA	U59524	Muerhoff et al., 1996
Isolate 8	2a	USA	U59525	Muerhoff et al., 1996
Isolate 9	2a	USA	U59526	Muerhoff et al., 1996
Isolate 10	2a	Europe	U59527	Muerhoff et al., 1996
Isolate 11	2a	Europe	U59528	Muerhoff et al., 1996
Isolate 12	2b	Europe	U59529	Muerhoff et al., 1996
Isolate 13	2b	Europe	U59530	Muerhoff et al., 1996
Isolate 14	2b	USA	U59531	Muerhoff et al., 1996
Isolate 15	2b	USA	U59532	Muerhoff et al., 1996
Isolate 16	2b	Europe	U59533	Muerhoff et al., 1996
Isolate 17	2b	Europe	U59534	Muerhoff et al., 1996
Isolate 18	2b	USA	U59535	Muerhoff et al., 1996
Isolate 19	2b	USA	U59536	Muerhoff et al., 1996
Isolate 20	2b	Europe	U59537	Muerhoff et al., 1996
Isolate 21	3	Asia	U59538	Muerhoff et al., 1996
Isolate 22	3	Asia	U59539	Muerhoff et al., 1996
K606	3	Japan	D87249	Fukushi et al., 1996, unpublished
K1737	3	Japan	D87250	Fukushi et al., 1996, unpublished
K1789	3	Japan	D87251	Fukushi et al., 1996, unpublished
K1916	3	Japan	D87252	Fukushi et al., 1996, unpublished
K2141	3	Japan	D87253	Fukushi et al., 1996, unpublished
K2142	3	Japan	D87254	Fukushi et al., 1996, unpublished

(62%) of the hemophiliacs were GBV-C RNA negative in serum by PCR (age range: 1–31; mean \pm SD: 12.7 \pm 7.6). HCV RNA was detected by nested-PCR using primers from the 5'UTR in 12 (27%) of the hemophiliacs (age range: 8–31; mean \pm SD: 16.5 \pm 6.2). Twenty-six (58%) of the hemophiliacs were anti-HCV positive (age range: 5 to 31; mean \pm SD: 14.4 \pm 6.1), while 19 (42%) were anti-HCV negative by EIA (age range: 1–23; mean \pm SD: 10.2 \pm 7.8). All the anti-HCV positive but HCV RNA negative samples were also tested by RIBA and confirmed positive. Dual infections with GBV-C and HCV were detected in 8 (18%) of the patients (age range: 8–20; mean \pm SD: 14.4 \pm 4.5), of whom 2 were anti-HCV positive but negative for HCV

TABLE II. Frequencies of HCV and GBV-C Markers in 45 Nicaraguan Hemophiliacs

Age range (yrs)	n	GBV-C RNA	HCV RNA	Anti-HCV	Any marker
1–10	19	7 (37%)	2 (10%)	8 (42%)	13 (68%)
11–20	19	9 (47%)	8 (42%)	14 (74%)	17 (89%)
21–>30	7	1 (14%)	2 (29%)	4 (57%)	5 (71%)
Total	45	17 (38%)	12 (27%)	26 (58%)	35 (78%)

RNA. Ten (22%) of the hemophiliacs were negative for both viruses (age range: 1–23; mean \pm SD: 9.6 \pm 8.5). HCV RNA was more often detected in sera of hemophiliacs above 10 years in age than in the younger age




Fig. 1. Dendrogram obtained with UPGMA based on the 5'UTR sequences from 11 Nicaraguan hemophiliacs and 49 GBV-C sequences retrieved from GeneBank and 3 sequences from GBV-C strains from Swedish drug addicts. The figures at the forks of the dendrogram indicate the bootstrap values obtained from 100 replicas.

group, 10 out of 26 versus 2 out of 19 ($P < 0.05$; Fisher's test). No such relation was found for the presence of GBV-C RNA or anti-HCV. The mean age of patients positive for GBV-C but negative for HCV by PCR was significantly lower than for those negative for GBV-C but positive for HCV by PCR, 11.0 versus 18.3 years, respectively ($P < 0.05$; Student's *t*-test). Thus GBV-C infections in the Nicaraguan hemophiliacs seem to occur earlier than the infections with HCV.

As shown in Figure 1, a phylogenetic analysis of 5'UTR in 11 of the 17 Nicaraguan GBV-C strains showed that they were most similar to Asian GBV-C strains, although they formed a separate cluster as

compared to the Asian strains provisionally designated as genotype 3 [Muerhoff et al., 1996].

Sequences from the NS5-B region were available in 8 of the 12 HCV strains recovered, all of which were found to belong to genotype 1a.

DISCUSSION

High prevalence rates for both GBV-C and HCV were observed in the group of hemophiliacs studied and in all age groups in agreement with previous studies reporting a high risk for HCV infection [Brettler et al., 1990; Rumi et al., 1990; Makris et al., 1990; Simmonds

Table III. Age Distribution of Nicaraguan Hemophiliacs According to Detection of GBV-C and HCV RNA by PCR

Characteristic	GBV-C and HCV RNA			
	GBV-C (+) HCV (-)	GBV-C (+) HCV (+)	GBV-C (-) HCV (+)	GBV-C (-) HCV (-)
Number of cases	11	6	6	22
Mean age \pm SD	11.0 \pm 6.8 ^a	14.7 \pm 5.3	18.3 \pm 7.0 ^a	11.1 \pm 7.1
Anti-HCV	2 (18%)	6 (100%)	6 (100%)	12 (54%)

^a*P* < 0.05; Student's *t*-test.

et al., 1990; Naoumov and Rumi, 1991] as well as an increased risk for GBV-C infection in patients with this background [Schleicher et al., 1996; Tagariello et al., 1996]. Dual infections have also been reported owing to both viruses sharing transmission pathways [Masuko et al., 1996; Nübling and Lower, 1996; Feucht et al., 1996; Linnen et al., 1996]. In this study a dual infection with GBV-C was detected in 8 (31%) of 26 HCV infected patients. All these patients were positive both by PCR and by a second-generation anti-HCV EIA. The lower age of the hemophiliacs with GBV-C infections as compared to those with HCV infections indicate that the risk for this group of patients to acquire GBV-C infection is higher than that for HCV infection. There was a high frequency of infections due to both GBV-C and HCV in the present study, considering the rather low prevalence of HCV, 0.5%, observed in donors from Nicaragua [Garcia et al., 1996].

Although the epidemiology of GBV-C remains to be clarified, recent studies in the United States and Japan have shown a relatively high frequency of GBV-C infections among blood donors, 1.7% and 0.8%, respectively [Dawson et al., 1996; Linnen et al., 1996; Masuko et al., 1996]. Transmission of the virus has been demonstrated to occur by blood transfusions and by other parenteral modes [Aikawa et al., 1996; Masuko et al., 1996; Schmidt et al., 1996] as well as vertically from mother to newborn child [Feucht et al., 1996]. The virus has been found in patients suffering from both acute and chronic cryptogenic hepatitis [Simons et al., 1995; Fiordalisi et al., 1996; Leary et al., 1996] and has also been implicated in fulminant hepatitis [Yoshida et al., 1995; Heringlake et al., 1996], although this outcome of GBV-C infection still remains controversial [Kuroki et al., 1996; Kao et al., 1996].

The finding of genotype 1a of HCV in Nicaragua was anticipated, since this genotype is prevalent in the Americas [Andonov and Chaudhary, 1994; Cha et al., 1992; Simmonds et al., 1993; Stuyver et al., 1993] but is also distributed worldwide [McOmish et al., 1994]. Some HCV genotypes (i.e., 4–11) are mainly confined to restricted areas in Africa and Asia [Cha et al., 1992; Simmonds et al., 1994; Mellor et al., 1995; Tokita et al., 1994a, 1994b, 1995]. However, none of these genotypes was found in the studied population. Therefore, GBV-C strains sharing sequence similarities to strains with Asian background was unexpected. A possible explanation for this finding would be that the GBV-C strains in Central America have their origins in Amerindian

populations, since all the hemophiliacs had been treated only with locally derived blood products and no commercial cryoprecipitates from the United States or Japan had been used. Molecular epidemiology may, apart from its use in tracing routes of viral transmission as for HBV [Magnius & Norder, 1995], HCV [Chou et al., 1993; Davidson et al., 1995], and GBV-C [Egawa et al., 1996; Schmidt et al., 1996], also be used as a tool to study prehistoric and historic migrations of human population as for, e.g., HTLV I [Yanagihara, 1994; Song et al., 1995] and HBV [Norder et al., 1993]. Recent phylogenetic studies on HTLV II sequences, as well as genetic studies, favour the idea that the ancestral migration waves preceeding the settlement of the New World have origins in Mongolia, Manchuria, or Southeast Siberia [Hall et al., 1994; Neel et al., 1994; Biggar et al., 1996]. That Amerindian viral strains may be prevalent in Central American populations with Hispanic background was recently described for HBV, where the prevailing genotype F strains also are the strains prevailing in the Amerindian populations of the Americas [Arauz-Ruiz et al., 1997]. The similarity of Nicaraguan GBV-C strains to strains from Asia, suggesting that these strains have Amerindian origin, indicates that GBV-C is an extremely conserved virus and explains why the divergence of GBV-C strains with different geographical origin is much lower than that for HCV. Future studies will establish whether the geographical variants of GBV-C, although most likely representing ancient splittings, diverge to the extent needed to define them as genotypes of GBV-C.

ACKNOWLEDGMENTS

We thank Noel McCarthy for correcting the English and Anna Eriksson for skillful technical assistance. This work was supported by research grants from KIRT, SAREC, and the Swedish Medical Research Council (K96-16P-11019-03A). The nucleotide sequence data reported in this paper will appear in the EMBL, GenBank, and DDBJ Nucleotide Sequence Databases under accession no. V86113-V86126.

REFERENCES

- Aikawa T, Sugai Y, Okamoto H (1996): Hepatitis G infection in drug abusers with chronic hepatitis C. *New England Journal of Medicine* 334:195–196.

- Andonov A, Chaudhary RK (1994): Genotyping of Canadian hepatitis C virus isolates by PCR. *Journal of Clinical Microbiology* 32:2031-2034.
- Arauz-Ruiz P, Norder H, Visoná KA, Magnius LO (1997): Genotype F prevails in HBV infected patients of Hispanic origin in Central America and may carry the precore stop mutant. *Journal of Medical Virology* 51:305-312.
- Biggar RJ, Taylor ME, Neel JV, Hjelle B, Levine PH, Black FL, Shaw GM, Sharp PM, Hahn BH (1996): Genetic variants of human T-lymphotropic virus type II in American Indian groups. *Virology* 216:165-173.
- Brettler DB, Alter HJ, Dienstag JL, Forsberg AD, Levine PH (1990): Prevalence of hepatitis C virus antibody in a cohort of hemophilia patients. *Blood* 76:254-256.
- Casanova JL, Penetier C, Jaulin C, Kourilsky P (1990): Optimal conditions for directly sequencing double-stranded PCR products with Sequenase. *Nucleic Acids Research* 18:4028.
- Casey JL, Brown TL, Colan EJ, Wignall FS, Gerin JL (1993): A genotype of hepatitis D virus that occurs in northern South America. *Proceedings of the National Academy of Sciences of the USA* 90:9016-9020.
- Cha TA, Beall E, Irvine B, Kolberg J, Chien D, Kuo G, Urdea MS (1992): At least five related, but distinct, hepatitis C viral genotypes exist. *Proceedings of the National Academy of Sciences of the USA* 89:7144-7148.
- Choo QL, Richman K, Han JH, Berger K, Lee C, Dong C, Gallegos C, Coit D, Medina-Selby A, Barr PJ, Weiner A, Bradley DW, Kuo G, Houghton M (1991): Genetic organization and diversity of the hepatitis C virus. *Proceedings of the National Academy of Sciences of the USA* 88:2451-2455.
- Chou WH, Lin SF, Sheu SH, Lu CF, Lin SY, Wu JS (1993): Comparison of hepatitis C virus strains obtained from hemodialysis patients. *Japanese Journal of Medical Science and Biology* 46:191-202.
- Davidson F, Simmonds P, Ferguson JC, Jarvis LM, Dow BC, Follett EAC, Seed CRG, Krusius T, Lin C, Medgyesi GA, Kiyokawa H, Olim G, Duraisamy G, Cuypers T, Saeed AA, Teo D, Conradie J, Kew MC, Lin M, Nuchaprayoon C, Ndimbie OK, Yap PL (1995): Survey of major genotypes and subtypes of hepatitis C virus using RFLP of sequences amplified from the 5' non-coding region. *Journal of General Virology* 76:1197-1204.
- Dawson GJ, Schlauder GG, Pilot-Matias TJ, Thiele D, Leary TP, Murphy P, Rosenblatt JE, Simons JN, Martinson FEA, Gutierrez RA, Lentino JR, Pachucki C, Muerhoff AS, Widell A, Tegtmeier G, Desai S, Mushahwar IK (1996): Prevalence studies of GB virus-C infection using reverse transcriptase-polymerase chain reaction. *Journal of Medical Virology* 50:97-103.
- Egawa K, Yukawa T, Arakawa S, Nakao H, Inoue T, Tanaka T, Tsuda F, Okamoto H, Miyakawa Y, Mayumi M (1996): Infection with GB virus C in leprosy patients in Japan. *Journal of Medical Virology* 49:110-114.
- Felsenstein J (1993): PHYLIP: Phylogeny inference package. Version 3.52c. Seattle: University of Washington.
- Feucht HH, Zöllner B, Polywka S, Laufs R (1996): Vertical transmission of hepatitis G. *Lancet* 347:615-616.
- Fiordalisi G, Zanella I, Mantero G, Bettinardi A, Stellini R, Paraninfo G, Cadeo G, Primi D (1996): High prevalence of GB virus C infection in a group of Italian patients with hepatitis of unknown etiology. *Journal of Infectious Diseases* 174:181-183.
- García Z, Taylor L, Ruano A, Pavón L, Ayerdis E, Luftig RB, Visoná KA (1996): Evaluation of a pooling method for routine anti-HCV screening of blood banks in countries under development. *Journal of Medical Virology* 49:218-222.
- Garson JA, Ring C, Tuke P, Tedder RS (1990): Enhanced detection by PCR of hepatitis C virus RNA. *Lancet* 336:878-879.
- Hall WW, Zhu SW, Horal P, Furuta Y, Zagaany G, Vahlne A (1994): HTLV-II infection in Mongolia. *AIDS Research and Human Retroviruses* 10:443.
- Heringlake S, Osterkamp S, Trautwein C, Tillmann HL, Böker K, Muerhoff S, Mushahwar IK, Hunsmann G, Manns MP (1996): Association between fulminant hepatic failure and a strain of GBV virus C. *Lancet* 348:1626-1629.
- Huang CC, Nguyen D, Fernandez J, Yun KY, Fry KE, Bradley DW, Tam AW, Reyes GR (1992): Molecular cloning and sequencing of the Mexico isolate of hepatitis E virus (HEV). *Virology* 191:550-558.
- Kao JH, Chen PJ, Chen DS (1996): GBV-C in the aetiology of fulminant hepatitis. *Lancet* 347:120.
- Kuroki T, Nishiguchi S, Tanaka M, Enomoto M, Kobayashi K (1996): Does GBV-C cause fulminant hepatitis in Japan? *Lancet* 347:908.
- Leary TP, Muerhoff AS, Simons JN, Pilot-Matias TJ, Erker JC, Chalmers ML, Schlauder GG, Dawson GJ, Desai SM, Mushahwar IK (1996): Consensus oligonucleotide primers for the detection of GB virus C in human cryptogenic hepatitis. *Journal of Virological Methods* 56:119-121.
- Leitner T, Halapi E, Scarlatti G, Rossi P, Albert J, Fenyo EM, Uhlén M (1993): Analysis of heterogeneous viral populations by direct DNA sequencing. *Bio Techniques* 15:120-127.
- Linnen J, Wages J, Zhang-Keck ZY, Fry KE, Krawczynski KZ, Alter H, Koonin E, Gallagher M, Alter M, Hadziyannis S, Karayiannis P, Fung K, Nakatsuji Y, Shih JWK, Young L, Piatak M, Hoover C, Fernandez J, Chen S, Zou JC, Morris T, Hyams KC, Ismay S, Lifson JD, Hess G, Fong SKH, Thomas H, Bradley D, Margolis H, Kim JP (1996): Molecular cloning and disease association of hepatitis G virus: A transfusion-transmissible agent. *Science* 271:505-508.
- Magnius LO, Norder H (1995): Subtypes, genotypes and molecular epidemiology of the hepatitis B virus as reflected by sequence variability of the S-gene. *Intervirology* 38:24-34.
- Makris M, Preston FE, Triger DR, Underwood JCE, Choo QL, Kuo G, Houghton M (1990): Hepatitis C antibody and chronic liver disease in haemophilia. *Lancet* 335:1117-1119.
- Masuko K, Mitsui T, Iwano K, Yamazaki C, Okuda K, Meguro T, Murayama N, Inoue T, Tsuda F, Okamoto H, Mirakawa Y, Mayumi M (1996): Infection with hepatitis GB virus C in patients on maintenance hemodialysis. *New England Journal of Medicine* 334:1485-1490.
- McOmish F, Yap PL, Dow BC, Follett EAC, Seed C, Keller AJ, Cobain TJ, Krusius T, Kolho E, Naukkarinen R, Lin C, Lai C, Leong S, Medgyesi GA, Héjjas M, Kiyokawa H, Fukada K, Cuypers T, Saeed AA, Al-Rasheed AM, Lin M, Simmonds P (1994): Geographical distribution of hepatitis C virus genotypes in blood donors: An international collaborative survey. *Journal of Clinical Microbiology* 32:884-892.
- Mellor J, Holmes EC, Jarvis LM, Yap PL, Simmonds P, and the International HCV Collaborative Study Group (1995): Investigation of the pattern of hepatitis C virus sequence diversity in different geographical regions: Implications for virus classification. *Journal of General Virology* 76:2493-2507.
- Muerhoff AS, Simons JN, Leary TP, Erker JC, Chalmers ML, Pilot-Matias TJ, Dawson GJ, Desai SM, Mushahwar IK (1996): Sequence heterogeneity within the 5'-terminal region of the hepatitis GB virus C genome and evidence for genotypes. *Journal of Hepatology* 25:379-384.
- Naoumov NV, Rumi MG (1991): Hepatitis C virus infection in hemophiliacs. *Annals of Internal Medicine* 114:171-172.
- Naumann H, Schaefer S, Yoshida CFT, Gaspar AMC, Repp R, Gerlich W (1993): Identification of a new hepatitis B virus (HBV) genotype from Brazil that expresses HBV surface antigen subtype adw4. *Journal of General Virology* 74:1627-1632.
- Neel JV, Biggar RJ, Sukernik RI (1994): Virologic and genetic studies relate Amerind origins to the indigenous people of the Mongolia/Manchuria/Southeastern Siberia region. *Proceedings of the National Academy of Sciences of the USA* 91:10737-10741.
- Norder H, Hammas B, Lee SD, Bile K, Couroucé AM, Mushahwar IK, Magnius LO (1993): Genetic relatedness of hepatitis B viral strains of diverse geographical origin and natural variations in the primary structure of the surface antigen. *Journal of General Virology* 74:1341-1348.
- Nübling CM, Löwer J (1996): GB-C genomes in a high-risk group, in plasma pools, and in intravenous immunoglobulin. *Lancet* 347:68.
- Okamoto H, Kurai K, Okada SI, Yamamoto K, Lizuka H, Tanaka T, Fukuda S, Tsuda F, Mishiro S (1992): Full-length sequence of a hepatitis C virus genome having poor homology to reported isolates: Comparative study of four distinct genotypes. *Virology* 188:331-341.
- Rumi MG, Colombo M, Gringeri A, Mannucci PM (1990): High prevalence of antibody to hepatitis C virus in multitransfused hemophiliacs with normal transaminase levels. *Annals of Internal Medicine* 112:379-380.
- Schleicher S, Chaves RL, Dehmer T, Gregor M, Hess G, Flehmig B (1996): Identification of GBV-C hepatitis G RNA in chronic hepatitis C patients. *Journal of Medical Virology* 50:71-74.

- Schmidt B, Korn K, Fleckenstein B (1996): Molecular evidence for transmission of hepatitis G virus by blood transfusion. *Lancet* 347: 909.
- Simmonds P, Zhang LQ, Watson HG, Rebus S, Ferguson ED, Balfe P, Leadbetter GH, Yap PL, Peutherer JF, Ludlam CA (1990): Hepatitis C quantification and sequencing in blood products, haemophiliacs, and drug users. *Lancet* 336:1469–1472.
- Simmonds P, McOmish F, Yap PL, Chan SW, Lin CK, Dusheiko G, Saeed AA, Holmes EC (1993): Sequence variability in the 5' non-coding region of hepatitis C virus: Identification of a new virus type and restrictions on sequence diversity. *Journal of General Virology* 74:661–668.
- Simmonds P, Alberti A, Alter HJ, Bonino F, Bradley DW, Brechot C, Brouwer JT, Chan SW, Chayama K, Chen DS, Choo QL, Colombo M, Cuypers HTM, Date T, Dusheiko GM, Esteban JI, Fay O, Hadziyannis SJ, Han J, Hatzakis A, Holmes EC, Hotta H, Houghton M, Irvine B, Kohata M, Kolberg JA, Kuo G, Lau JYN, Lelie PN, Maertens G, McOmish F, Miyamura T, Mizokami M, Nomoto A, Prince AM, Reesink HW, Rice C, Roggendorf M, Schalm SW, Shikata T, Shimotohno K, Stuyver L, Trépo C, Weiner A, Yap PL, Urdea MS (1994): A proposed system for the nomenclature of hepatitis C viral genotypes. *Hepatology* 19:1321–1324.
- Simons JN, Leary TP, Dawson GJ, Pilot-Matias TJ, Muerhoff AS, Schlauder GG, Desai SM, Mushahwar IK (1995): Isolation of novel virus-like sequences associated with human hepatitis. *Nature Medicine* 1:564–569.
- Song K, Nerurkar VR, Pereira-Cortez AJ, Yamamoto M, Taguchi H, Miyoshi I, Yanagihara R (1995): Sequence and phylogenetic analyses of human T cell lymphotropic virus type I from a Brazilian woman with adult T cell leukemia: Comparison with virus strains from South America and the Caribbean basin. *American Journal of Tropical Medicine and Hygiene* 52:101–108.
- Stuyver L, Rossau R, Wyseur A, Duhamel M, Vanderborght B, Van Heuverswyn H, Maertens G (1993): Typing of hepatitis C virus isolates and characterization of new subtypes using a line probe assay. *Journal of General Virology* 74:1093–1102.
- Tagariello G, Infantolino D, Biasin MR, Davoli PG, Traldi A (1996): Hepatitis G viral RNA in Italian haemophiliacs with and without hepatitis C infection. *Lancet* 348:760–761.
- Tokita H, Shrestha SM, Okamoto H, Sakamoto M, Horikata M, Iizuka H, Shrestha S, Miyakawa Y, Mayumi M (1994a): Hepatitis C virus variants from Nepal with novel genotypes and their classification into the third major group. *Journal of General Virology* 75:931–936.
- Tokita H, Okamoto H, Tsuda F, Song P, Nakata S, Chosa T, Iizuka H, Mishiho S, Miyakawa Y, Mayumi M (1994b): Hepatitis C virus variants from Vietnam are classifiable into the seventh, eighth, and ninth major genetic groups. *Proceedings of the National Academy of Sciences of USA* 91:11022–11026.
- Tokita H, Okamoto H, Luengrojanakul P, Vareesangthip K, Chainuvat T, Iizuka H, Tsuda F, Miyakawa Y, Mayumi M (1995): Hepatitis C virus variants from Thailand classifiable into five novel genotypes in the sixth (6b), seventh (7c,7d) and ninth (9b,9c) major genetic groups. *Journal of General Virology* 76:2329–2335.
- Yanagihara R (1994): Geographic-specific genotypes or topotypes of human T-cell lymphotropic virus type I as markers for early and recent migrations of human populations. *Advances in Virus Research* 43:147–186.
- Yoshida M, Okamoto H, Mishiho S (1995): Detection of the GBV-C hepatitis virus genome in serum from patients with fulminant hepatitis of unknown aetiology. *Lancet* 346:1131–1132.
- Yun Z, Lara C, Johansson B, Lorenzana de Rivera I, Sönnnerborg A (1996): Discrepancy of hepatitis C virus genotypes as determined by phylogenetic analysis of partial NS5 and core sequences. *Journal of Medical Virology* 49:155–160.
- Zuckerman AJ (1996): Alphabet of hepatitis viruses. *Lancet* 347:558–559.